

Federal Employee Program.

Federal Employee Program® 750 9th St NW Washington, D.C. 20001 202.942.1000 Fax 202.942.1125

5.90.055

Section: Prescription Drugs Effective Date: April 1, 2024

Subsection: Topical Products Original Policy Date: April 29, 2022

Subject: Cibingo Page: 1 of 7

Last Review Date: March 8, 2024

Cibingo

Description

Cibingo (abrocitinib)

Background

Cibinqo (abrocitinib) is a Janus kinase (JAK) inhibitor. JAKs are intracellular enzymes which transmit signals arising from cytokine or growth factor-receptor interactions on the cellular membrane to influence cellular processes of hematopoiesis and immune cell function. Within the signaling pathway, JAKs phosphorylate and activate Signal Transducers and Activators of Transcription (STATs) which modulate intracellular activity including gene expression. Cibinqo modulates the signaling pathway at the point of JAKs, preventing the phosphorylation and activation of STATs (1).

Regulatory status

FDA-approved indication: Cibinqo is a Janus kinase (JAK) inhibitor indicated for the treatment of adults and pediatric patients 12 years of age and older with refractory, moderate-to-severe atopic dermatitis whose disease is not adequately controlled with other systemic drug products, including biologics, or when use of those therapies are inadvisable (1).

Limitations of Use: (1)

Cibinqo is not recommended for use in combination with other JAK inhibitors, biologic immunomodulators, or with other immunosuppressants.

Cibingo carries several boxed warnings: (1)

Subsection: Topical Products Original Policy Date: April 29, 2022

Subject: Cibinqo Page: 2 of 7

1. Serious infections

a. Serious infections, including tuberculosis and bacterial, invasive fungal, viral, and other opportunistic infection leading to hospitalization or death. If a serious infection develops, discontinue Cibinqo until the infection is controlled. Prior to starting Cibinqo, perform a test for latent tuberculosis; if it is positive, start treatment for tuberculosis prior to starting Cibinqo. Monitor all patients for active tuberculosis during treatment, even if the initial latent tuberculosis test is negative.

2. Mortality

- a. RA patients 50 years of age and older with at least one cardiovascular risk factor showed a higher rate of all-cause mortality, including sudden cardiovascular death, in patients treated with JAK inhibitors compared to TNF blockers.
- b. Cibingo is not approved for use in RA patients.

3. Malignancies

- a. Lymphoma and other malignancies have been observed in patients treated with Cibingo.
- b. In RA patients treated with a JAK inhibitor, a higher rate of malignancies was observed when compared with TNF blockers.
- 4. Major adverse cardiovascular events (MACE)
 - a. RA patients 50 years of age and older with at least one cardiovascular risk factor treated with a JAK inhibitor showed a higher rate of MACE (defined as cardiovascular death, myocardial infarction, and stroke) when compared to TNF blockers. Patients who are current or past smokers are at increased risk. Cibinqo should be discontinued in patients that have experienced a myocardial infarction or stroke.

5. Thrombosis

- a. Thrombosis, including deep vein thrombosis, pulmonary embolism, and arterial thrombosis, have occurred in patients treated with JAK inhibitors used to treat inflammatory conditions. Many of these adverse events were serious and some resulted in death.
- b. In RA patients 50 years of age and older with at least one cardiovascular risk factor treated with a JAK inhibitor, a higher rate of thrombosis was observed when compared with TNF blockers.

Cibinqo is contraindicated in patients taking antiplatelet therapies, except for low-dose aspirin (≤81 mg daily), during the first 3 months of treatment (1).

Subsection: Topical Products Original Policy Date: April 29, 2022

Subject: Cibingo Page: 3 of 7

The safety and effectiveness of Cibinqo in pediatric patients less than 12 years of age have not been established (1).

Related policies

Adbry, Dupixent, Rinvoq

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Cibingo may be considered **medically necessary** if the conditions indicated below are met.

Cibingo may be considered investigational for all other indications.

Prior-Approval Requirements

Age 12 years of age or older

Diagnosis

Patient must have the following:

1. Moderate-to-severe atopic dermatitis (eczema)

AND ALL of the following:

- 1. Inadequate treatment response, intolerance, or contraindication to at least **TWO** systemic atopic dermatitis medications, including biologics (e.g., oral corticosteroids, hydroxyzine, Adbry, Dupixent, Rinvoq, etc.)
- 2. Prescriber has considered the risks for malignancy and major adverse cardiovascular events (MACE) (e.g., advanced age, smoking history, cardiovascular risk factors etc.) and determined that Cibinqo therapy is appropriate
- 3. Result for latent TB infection is negative **OR** result was positive for latent TB and patient completed treatment (or is receiving treatment) for latent TB

AND NONE of the following:

Subsection: Topical Products Original Policy Date: April 29, 2022

Subject: Cibinqo Page: 4 of 7

1. Antiplatelet therapy (excluding low-dose aspirin ≤81 mg daily) during the first 3 months of treatment

- 2. Active bacterial, invasive fungal, viral, and other opportunistic infections
- 3. Severe hepatic impairment (Child Pugh C)
- 4. A lymphocyte count less than 500 cells/mm3
- 5. An absolute neutrophil count less than 1000 cells/mm3
- 6. A hemoglobin less than 8 g/dL
- 7. History of thrombotic events including deep vein thrombosis (DVT) or pulmonary embolism (PE)
- 8. Used in combination with another non-topical Prior Authorization (PA) medication for atopic dermatitis (see Appendix 1)
- 9. Given concurrently with live vaccines

Prior-Approval Renewal Requirements

Age 12 years of age or older

Diagnosis

Patient must have the following:

1. Atopic dermatitis (eczema)

AND ALL of the following:

- 1. Condition has improved or stabilized
- 2. Prescriber has considered the risks for malignancy and major adverse cardiovascular events (MACE) (e.g., advanced age, smoking history, cardiovascular risk factors etc.) and determined that continuation of Cibinqo therapy is appropriate

AND NONE of the following:

- 1. Active bacterial, invasive fungal, viral, and other opportunistic infections
- 2. Used in combination with another non-topical Prior Authorization (PA) medication for atopic dermatitis (see Appendix 1)
- 3. Development of thrombotic events (including DVTs or PEs)
- 4. Given concurrently with live vaccines

Subsection: Topical Products Original Policy Date: April 29, 2022

Subject: Cibinqo Page: 5 of 7

Policy Guidelines

Pre-PA Allowance

None

Prior-Approval Limits

Quantity 90 tablets per 90 days

Duration 4 months

Prior-Approval Renewal Limits

Quantity 90 tablets per 90 days

Duration 12 months

Rationale

Summary

Cibinqo (abrocitinib) is a Janus kinase (JAK) inhibitor indicated for patients with atopic dermatitis. Cibinqo has several boxed warnings including risk of serious infections, mortality, malignancies, MACE, and thrombosis. The safety and effectiveness of Cibinqo in pediatric patients less than 12 years of age have not been established (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Cibingo while maintaining optimal therapeutic outcomes.

References

1. Cibinqo [package insert]. New York, NY: Pfizer Inc.; December 2023.

Policy History

Date Action

April 2022 Addition to PA

5.90.055

Section:Prescription DrugsEffective Date:April 1, 2024Subsection:Topical ProductsOriginal Policy Date:April 29, 2022

Subject: Cibinqo Page: 6 of 7

June 2022 Annual review. Per SME, added example names of biologics to the atopic

dermatitis t/f requirement

March 2023 Per PI update, changed age requirement to 12 and older from 18 and

older. Changed policy number to 5.90.055

June 2023 Annual review

March 2024 Annual review and reference update

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 8, 2024 and is effective on April 1, 2024.

5.90.055

Section: Prescription Drugs Effective Date: April 1, 2024

Subsection: Topical Products Original Policy Date: April 29, 2022

Subject: Cibinqo Page: 7 of 7

Appendix 1 - List of Non-Topical PA Medications for Atopic Dermatitis

Generic Name	Brand Name
abrocitinib	Cibinqo
dupilumab	Dupixent
tralokinumab-ldrm	Adbry
upadactinib	Rinvoq