



FEP Medical Policy Manual

FEP 7.01.95 Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Annual Effective Policy Date: January 1, 2025

Original Policy Date: June 2012

Related Policies:

- 6.01.10 - Stereotactic Radiosurgery and Stereotactic Body Radiotherapy
- 7.01.91 - Radiofrequency Ablation of Primary or Metastatic Liver Tumors
- 7.01.92 - Cryoablation of Tumors Located in the Kidney, Lung, Breast, Pancreas, or Bone

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Description

Description

In radiofrequency ablation (RFA), a probe is inserted into the center of a tumor; then, prong-shaped, non-insulated electrodes are projected into the tumor. Next, heat is generated locally by an alternating, high-frequency current that travels through the electrodes. The localized heat treats the tissue adjacent to the probe, resulting in a 3 cm to 5.5 cm sphere of dead tissue. The cells killed by RFA are not removed but are gradually replaced by fibrosis and scar tissue. If there is a local recurrence, it occurs at the edge and can sometimes be retreated. RFA may be performed percutaneously, laparoscopically, or as an open procedure.

OBJECTIVE

The objective of this evidence review is to determine whether the use of radiofrequency ablation improves the net health outcome in individuals with a range of tumors (including osteolytic bone metastases, osteoid osteomas, renal cell carcinoma, lung cancer, breast tumors, thyroid tumors, and others).

POLICY STATEMENT

Osteolytic bone metastases

Radiofrequency ablation may be considered **medically necessary** to palliate pain in individuals with osteolytic bone metastases who have failed or are poor candidates for standard treatments such as radiation or opioids.

Osteoid osteomas

Radiofrequency ablation may be considered **medically necessary** to treat osteoid osteomas that cannot be managed successfully with medical treatment.

Renal cell carcinoma

Radiofrequency ablation may be considered **medically necessary** to treat localized renal cell carcinoma that is no more than 4 cm in size when criteria 1 and 2 are met:

1. When it is necessary to preserve kidney function in individuals with significantly impaired renal function (ie, the individual has 1 kidney or renal insufficiency defined by a glomerular filtration rate of <60 mL/min/m²);
2. When the standard surgical approach (ie, resection of renal tissue) is likely to worsen existing kidney function substantially; OR When the individual is not considered a surgical candidate.

Non-small-cell lung cancer

Radiofrequency ablation may be considered **medically necessary** to treat an isolated peripheral non-small-cell lung cancer lesion that is no more than 3 cm in size when criteria 1 and 2 are met:

1. When surgical resection or radiotherapy with curative intent is considered appropriate based on stage of disease, however, medical comorbidity renders the individual unfit for those interventions;
2. When the tumor is located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery, and the heart.

Nonpulmonary tumor(s) metastatic to the lung

Radiofrequency ablation may be considered **medically necessary** to treat malignant nonpulmonary tumor(s) metastatic to the lung that are no more than 3 cm in size when criteria 1 and 2 are met:

1. When it is necessary to preserve lung function because surgical resection or radiotherapy is likely to worsen pulmonary status substantially; OR When the individual is not considered a surgical candidate;
2. When there is no evidence of extrapulmonary metastases; AND the tumor is located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery, and the heart.

(See the Policy Guidelines section for additional criteria.)

Radiofrequency ablation is considered **investigational** as a technique for ablation of:

- breast tumors;
- lung cancer not meeting the criteria above;
- renal cell cancer not meeting the criteria above;
- osteoid osteomas that can be managed with medical treatment;
- painful bony metastases as initial treatment; and
- all other tumors outside the liver including, but not limited to, the head and neck, thyroid, pancreas, adrenal gland, ovary, and pelvic/abdominal metastases of unspecified origin.

POLICY GUIDELINES

The following are additional criteria developed by clinical judgment or consensus and existing guidelines for the use of radiofrequency ablation to treat metastatic tumors to the lung:

- No more than 3 tumors per lung should be ablated;
- Tumors should be amenable to complete ablation; AND
- Twelve months should elapse before a repeat ablation is considered.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

The U.S. Food and Drug Administration (FDA) issued a statement in September 2008, concerning the regulatory status of RFA. The FDA has cleared RFA devices for the general indication of soft tissue cutting, coagulation, and ablation by thermal coagulation necrosis. Under this general indication, RFA can be used to ablate tumors, including lung tumors. Some RFA devices have been cleared for additional specific treatment indications, including partial or complete ablation of nonresectable liver lesions and palliation of pain associated with metastatic lesions involving bone. The FDA has not cleared any RFA devices for the specific treatment indication of partial or complete ablation of lung tumors, citing lack of sufficient clinical data to establish safety and effectiveness for this purpose. The FDA has received reports of death and serious injuries associated with the use of RFA devices in the treatment of lung tumors.

RATIONALE

Summary of Evidence

For individuals who have painful osteolytic bone metastases who have failed or are poor candidates for standard treatments who receive radiofrequency ablation (RFA), the evidence includes a prospective cohort study and case series. Relevant outcomes are symptoms, change in disease status, quality of life (QOL), medication use, and treatment-related morbidity. A prospective cohort study and case series have shown clinically significant pain relief (defined as a decrease of 2 units from baseline on the Brief Pain Inventory scale) or reduction in opioid use following treatment of painful osteolytic metastases. A multicenter, prospective study reported significant reductions in pain through the 6-month follow-up period, with 59% of patients achieving immediate improvement in pain within 3 days of RFA. The population is comprised of patients with few or no treatment options, for whom short-term pain relief is an appropriate clinical outcome. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have painful osteoid osteomas who receive RFA, the evidence includes numerous observational studies and systematic reviews of these studies. Relevant outcomes are symptoms, change in disease status, QOL, medication use, and treatment-related morbidity. In a systematic review of thermal ablation techniques, clinical success (pain-free) was achieved in 94% to 98% of patients. Most patients (89% to 96%) remained pain-free when assessed during longer-term follow-up. Another systematic review reported similar success rates noting an average 8.3% failure rate among patients receiving computed tomography (CT)-guided RFA. Although no randomized trials of RFA for osteoid osteomas have been performed, the uncontrolled studies have demonstrated RFA can provide adequate symptom relief with minimal complications, for a population for whom short-term symptom relief and avoidance of invasive procedures are appropriate clinical outcomes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have localized renal cell carcinoma (RCC) that is no more than 4 cm in size who receive RFA, the evidence includes a randomized controlled trial (RCT), numerous observational studies, and systematic reviews of these studies. Relevant outcomes are overall survival (OS), change in disease status, QOL, and treatment-related morbidity. A recent meta-analysis that included only an RCT and cohort studies found that RFA was as effective as nephrectomy for small renal tumors, with a reduction in complications. Another recent meta-analysis found that partial nephrectomy (PN) was superior to ablative techniques (the study included RFA but also cryoablation and microwave ablation) in overall mortality and local recurrence but not in cancer-specific mortality. It also found fewer complications and improved renal function with ablation. A meta-analysis from 2022 found that PN was superior to ablation (RFA, cryoablation, and microwave ablation) in local recurrence. Overall complications, decline in renal function, and cancer-specific mortality rates did not differ between ablation and PN. Although inconsistent, the evidence does suggest that, for small renal tumors, RFA may result in a similar rate of disease progression with a lower complication rate than nephrectomy. However, comparative trials are needed to determine with greater certainty the effects of these treatments in the same patient population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have inoperable primary pulmonary tumors or nonpulmonary tumors metastatic to the lung who receive RFA, the evidence includes prospective observational studies and systematic reviews of these studies. Relevant outcomes are OS, change in disease status, QOL, and treatment-related morbidity. A multicenter study found that for tumors less than 3.5 cm in size, RFA can lead to a complete response in as many as 88% of patients for at least 1 year. Two-year survival rates have been reported to range from 41% to 75% in case series, with 5-year survival rates of 20% to 27%. In general, the evidence suggests that RFA results in adequate survival and tumor control in patients who are not surgical candidates, with low morbidity rates. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have breast tumors who receive RFA, the evidence includes observational studies and systematic reviews of these studies. Relevant outcomes are OS, change in disease status, QOL, and treatment-related morbidity. Evidence has reported varied and incomplete ablation rates with concerns about postablation tumor cell viability. Long-term improvements in health outcomes have not been demonstrated. Additionally, available studies do not permit comparisons with conventional breast-conserving procedures. Further prospective studies, with long-term follow-up, should focus on whether RFA of the breast for small tumors can provide local control and survival rates compared with conventional breast-conserving treatment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have benign thyroid tumors who receive RFA, the evidence includes RCTs, prospective studies, case series, and systematic reviews of these studies. Relevant outcomes are symptoms, change in disease status, QOL, medication use, and treatment-related morbidity. Systematic reviews have demonstrated that RFA results in a significant reduction in thyroid nodule size, with a 2020 review showing that these changes remain durable through at least 36 months and a 2024 review indicating durability up to 5 years. Complication rates are generally low but include voice changes. The data are limited by significant heterogeneity in meta-analyses, a lack of generalizability to populations outside Republic of Korea and Italy, and a lack of comparators more relevant to practice in the United States. Further studies comparing RFA to percutaneous ethanol injection (PEI) or surgery would be more informative in determining the potential utility of RFA in patients with symptomatic or large benign thyroid tumors as these are the recommended treatment options per the American Thyroid Association. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have miscellaneous tumors (eg, head and neck, thyroid cancer, pancreas) who receive RFA, the evidence includes a few case series, prospective observational studies, and retrospective comparative studies. Relevant outcomes are OS, change in disease status, QOL, and treatment-related morbidity. There is a limited evidence base for these tumor types. Reporting on outcomes or comparisons with other treatments is limited. These studies do not permit conclusions on the health benefits of RFA. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Chest Physicians

The American College of Chest Physicians (2013) guidelines on the treatment of stage I and II non-small-cell lung cancer (NSCLC) have indicated RFA has been used effectively in clinical stage I NSCLC.⁷³ Therefore, in medically inoperable patients, peripheral NSCLC tumors less than 3 cm may be treated with RFA. The College also collaborated with the Society of Thoracic Surgeons to develop consensus guidelines on the treatment of high-risk patients with stage I NSCLC.⁷⁴ These 2012 consensus guidelines indicated RFA is an alternative treatment option for patients who are not surgical candidates due to severe medical comorbidity.

American Head and Neck Society - Endocrine Surgery Section

An international, multidisciplinary consensus statement on RFA and related ultrasound-guided ablation technologies for the treatment of benign and malignant thyroid disease was released in 2022 through a collaboration of international professional societies, including the Endocrine Surgery Section of the American Head and Neck Society.⁷⁵ Select relevant recommendations from the guideline are listed in Table 1.

Table 1. Summary of RFA Recommendations for Treatment of Benign and Malignant Thyroid Disease*

| | |
|-------------------|---|
| Recommendation 1 | US-guided ablation procedures may be used as a first-line alternative to surgery for patients with benign thyroid nodules contributing to compressive and/or cosmetic symptoms. |
| Recommendation 2 | Although less efficacious than surgery or RAI in normalizing thyroid function, thermal ablation procedures can be a safe therapeutic alternative in patients with an autonomously functional thyroid nodule and contraindications to first-line techniques. |
| Recommendation 3a | US-guided ablation procedures may be considered in patients with suitable primary papillary microcarcinoma who are unfit for surgery or decline surgery or active surveillance |
| Recommendation 3b | US-guided ablation procedures may be considered in patients with suitable recurrent papillary thyroid carcinoma who are unfit for surgery or decline surgery or active surveillance |
| Recommendation 3c | Repeat ablation of a benign nodule can be considered for remnant nodular tissue contributing to unresolved symptomatic or cosmetic concerns |

*This is not a comprehensive list of recommendations from the guideline.

RAI: radioactive iodine; RFA: radiofrequency ablation; US: Ultrasound.

American Urological Association

The American Urological Association (AUA; 2017) guideline on renal masses and localized renal cancer affirms that partial nephrectomy should be prioritized for the management of cT1a renal masses when intervention is indicated.⁷⁶ Thermal ablation should be considered "as an alternate approach for the management of cT1a renal masses <3 cm in size." The guidelines were updated in 2021 and recommendations are generally consistent with what was published in the 2017 guideline.⁷⁷ The 2021 AUA guideline explicitly states that RFA and cryoablation may be offered as options to patients who elect thermal ablation.

American Thyroid Association

The American Thyroid Association (2015) guideline on the management of thyroid nodules and differentiated thyroid cancer provides recommendations for management.⁴⁹ Patients with a benign cytology diagnosis or those very unlikely to be malignant (eg, purely cystic nodule) should undergo surveillance with the frequency determined by the level of suspicion for a missed malignancy. Medical or surgical intervention is considered if the nodules are large (>4 cm), causing compressive or structural symptoms, or if there is clinical concern. Recurrent cystic thyroid nodules with benign cytology should be considered for surgical removal or percutaneous ethanol injection. For differentiated thyroid cancer, "localized treatments with thermal (radiofrequency or cryo-) ablation, ethanol ablation, or chemoembolization may be beneficial in patients with a single or a few metastases and in those with metastases at high risk of local complications."

National Comprehensive Cancer Network

The National Comprehensive Cancer Network (NCCN) guidelines for the treatment of NSCLC (v.2.2024) state:⁷⁸ "For medically operable disease, resection is the preferred local treatment modality (other modalities include SABR [stereotactic ablative radiotherapy], thermal ablation such as radiofrequency ablation, and cryotherapy)." For patients who are not amenable to surgery, image-guided thermal ablation therapy (IGTA; includes RFA, microwave ablation, and cryoablation) may be considered. The guidance states "IGTA is an option for the management of NSCLC lesions <3 cm. Ablation for NSCLC lesions >3 cm may be associated with higher rates of local recurrence and complications."

The NCCN guidelines for thyroid carcinoma (v.3.2024) indicate that local therapies such as RFA may be considered for locoregional recurrence of thyroid carcinoma-papillary carcinoma in select patients with limited burden nodal disease. Additionally, local therapies, including RFA, can be

considered in those with metastatic disease.⁷⁹

The NCCN guidelines (v.1.2025) for renal cell carcinoma indicate that "thermal ablation (eg, cryosurgery, radiofrequency ablation, microwave ablation) is an option for the management of clinical stage T1 renal lesions. Thermal ablation is an option for clinical T1b masses in select patients not eligible for surgery. Biopsy of lesions is recommended to be done prior to or at time of ablation. Ablative techniques may require multiple treatments to achieve the same oncologic outcomes as conventional surgery."⁸⁰

The NCCN colon cancer guidelines (v.4.2024) state that "resection is the standard approach for the local treatment of resectable metastatic disease. However, patients with liver or lung oligometastases can also be considered for tumor ablation therapy, particularly in cases that may not be optimal for resection."⁸¹ "There is extensive evidence on the use of RFA as a reasonable treatment option for non-surgical candidates and for recurrent disease after hepatectomy with small liver metastases that can be treated with clear margins."

The NCCN guidelines for head and neck cancers (v.4.2024),⁸² breast cancer (v.4.2024),⁸³ bone cancer (v.2.2024),⁸⁴ and pancreatic adenocarcinoma (v.2.2024) do not mention RFA.⁸⁵

National Institute for Health and Care Excellence

The NICE guidance (2004) on osteoid osteoma indicated that "current evidence on the safety and efficacy of computed tomography (CT)-guided thermocoagulation of osteoid osteoma appears adequate to support its use...."⁸⁶

Updated NICE guidance (2010) on renal cancer has indicated that "evidence on the safety and efficacy of percutaneous radiofrequency ablation (RFA) ... in the short and medium term appears adequate to support the use of this procedure provided that patients are followed up in the long term."⁸⁷

The NICE guidance (2010) on RFA for primary and secondary lung cancers has stated: "Current evidence on the efficacy of percutaneous radiofrequency ablation (RFA) ... is adequate in terms of tumor control."⁸⁸ The NICE also indicated RFA might "be used in patients with small, early-stage lung cancers or small numbers of lung metastases who are unsuitable for, or prefer not to undergo, surgery. It may also have a place in multi-modality treatment of more advanced primary lung cancers." The guidance warned of serious complications (eg, pneumothorax) among lung cancer patients.

The NICE guidance (2016) on benign thyroid nodules stated: "Current evidence on the safety and efficacy of ultrasound-guided percutaneous radiofrequency ablation ... is adequate to support the use of this procedure...."⁸⁹

Society of Interventional Radiology

The Society of Interventional Radiology (2020) published a position statement on the role of percutaneous ablation in renal cell carcinoma.⁹⁰ Their relevant recommendations are as follows:

- "In patients with small renal tumors (stage T1a), percutaneous thermal ablation is a safe and effective treatment with fewer complications than nephrectomy and acceptable long-term oncological and survival outcomes. (Level of Evidence: C; Strength of Recommendation: Moderate)"
- "In selected patients with suspected T1a renal cell carcinoma, percutaneous thermal ablation should be offered over active surveillance. (Level of Evidence: C; Strength of Recommendation: Moderate)"
- "In high-risk patients with T1b renal cell carcinoma who are not surgical candidates, percutaneous thermal ablation may be an appropriate treatment option; however, further research in this area is required. (Level of Evidence: D; Strength of Recommendation: Weak)"
- "Radiofrequency ablation, cryoablation, and microwave ablation are all appropriate modalities for thermal ablation, and method of ablation should be left to the discretion of the operating physician. (Level of Evidence: D; Strength of Recommendation: Weak)"

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Jatoi I, Sung H, Jemal A. The Emergence of the Racial Disparity in U.S. Breast-Cancer Mortality. *N Engl J Med*. Jun 23 2022; 386(25): 2349-2352. PMID 35713541
2. Yedjou CG, Sims JN, Miele L, et al. Health and Racial Disparity in Breast Cancer. *Adv Exp Med Biol*. 2019; 1152: 31-49. PMID 31456178
3. National Cancer Institute. SEER Cancer Stat Facts: Kidney and Renal Pelvis Cancer. 2024. <https://seer.cancer.gov/statfacts/html/kidrp.html>. Accessed July 29, 2024.
4. Howard JM, Nandy K, Woldu SL, et al. Demographic Factors Associated With Non-Guideline-Based Treatment of Kidney Cancer in the United States. *JAMA Netw Open*. Jun 01 2021; 4(6): e2112813. PMID 34106265
5. Levy J, Hopkins T, Morris J, et al. Radiofrequency Ablation for the Palliative Treatment of Bone Metastases: Outcomes from the Multicenter OsteoCool Tumor Ablation Post-Market Study (OPuS One Study) in 100 Patients. *J Vasc Interv Radiol*. Nov 2020; 31(11): 1745-1752. PMID 33129427
6. Goetz MP, Callstrom MR, Charboneau JW, et al. Percutaneous image-guided radiofrequency ablation of painful metastases involving bone: a multicenter study. *J Clin Oncol*. Jan 15 2004; 22(2): 300-6. PMID 14722039
7. Grnemyer DH, Schirp S, Gevargez A. Image-guided radiofrequency ablation of spinal tumors: preliminary experience with an expandable array electrode. *Cancer J*. 2002; 8(1): 33-9. PMID 11898806
8. Kojima H, Tanigawa N, Kariya S, et al. Clinical assessment of percutaneous radiofrequency ablation for painful metastatic bone tumors. *Cardiovasc Intervent Radiol*. 2006; 29(6): 1022-6. PMID 16988875
9. Tordjiman M, Perronne L, Madelin G, et al. CT-guided radiofrequency ablation for osteoid osteomas: a systematic review. *Eur Radiol*. Nov 2020; 30(11): 5952-5963. PMID 32518986
10. Lanza E, Thouvenin Y, Viala P, et al. Osteoid osteoma treated by percutaneous thermal ablation: when do we fail? A systematic review and guidelines for future reporting. *Cardiovasc Intervent Radiol*. Dec 2014; 37(6): 1530-9. PMID 24337349
11. Albisinni U, Facchini G, Spinnato P, et al. Spinal osteoid osteoma: efficacy and safety of radiofrequency ablation. *Skeletal Radiol*. Aug 2017; 46(8): 1087-1094. PMID 28497160
12. Lassalle L, Campagna R, Corcos G, et al. Therapeutic outcome of CT-guided radiofrequency ablation in patients with osteoid osteoma. *Skeletal Radiol*. Jul 2017; 46(7): 949-956. PMID 28429047
13. Rimondi E, Mavrogenis AF, Rossi G, et al. Radiofrequency ablation for non-spinal osteoid osteomas in 557 patients. *Eur Radiol*. Jan 2012; 22(1): 181-8. PMID 21842430
14. Sahin C, Oc Y, Ediz N, et al. The safety and the efficacy of computed tomography guided percutaneous radiofrequency ablation of osteoid osteoma. *Acta Orthop Traumatol Turc*. Sep 2019; 53(5): 360-365. PMID 31371131
15. Knudsen M, Riishede A, Lcke A, et al. Computed tomography-guided radiofrequency ablation is a safe and effective treatment of osteoid osteoma located outside the spine. *Dan Med J*. May 2015; 62(5). PMID 26050823
16. Rosenthal DI, Hornicek FJ, Torriani M, et al. Osteoid osteoma: percutaneous treatment with radiofrequency energy. *Radiology*. Oct 2003; 229(1): 171-5. PMID 12944597
17. Yanagisawa T, Mori K, Kawada T, et al. Differential efficacy of ablation therapy versus partial nephrectomy between clinical T1a and T1b renal tumors: A systematic review and meta-analysis. *Urol Oncol*. Jul 2022; 40(7): 315-330. PMID 35562311
18. Uhlig J, Strauss A, Rcker G, et al. Partial nephrectomy versus ablative techniques for small renal masses: a systematic review and network meta-analysis. *Eur Radiol*. Mar 2019; 29(3): 1293-1307. PMID 30255245
19. Katsanos K, Mailli L, Krokidis M, et al. Systematic review and meta-analysis of thermal ablation versus surgical nephrectomy for small renal tumours. *Cardiovasc Intervent Radiol*. Apr 2014; 37(2): 427-37. PMID 24482030
20. El Dib R, Touma NJ, Kapoor A. Cryoablation vs radiofrequency ablation for the treatment of renal cell carcinoma: a meta-analysis of case series studies. *BJU Int*. Aug 2012; 110(4): 510-6. PMID 22304329
21. Liu SY, Chu CM, Kong AP, et al. Radiofrequency ablation compared with laparoscopic adrenalectomy for aldosterone-producing adenoma. *Br J Surg*. Oct 2016; 103(11): 1476-86. PMID 27511444
22. Marshall HR, Shakeri S, Hosseiny M, et al. Long-Term Survival after Percutaneous Radiofrequency Ablation of Pathologically Proven Renal Cell Carcinoma in 100 Patients. *J Vasc Interv Radiol*. Jan 2020; 31(1): 15-24. PMID 31767409
23. Andrews JR, Atwell T, Schmit G, et al. Oncologic Outcomes Following Partial Nephrectomy and Percutaneous Ablation for cT1 Renal Masses. *Eur Urol*. Aug 2019; 76(2): 244-251. PMID 31060824
24. Park BK, Gong IH, Kang MY, et al. RFA versus robotic partial nephrectomy for T1a renal cell carcinoma: a propensity score-matched comparison of mid-term outcome. *Eur Radiol*. Jul 2018; 28(7): 2979-2985. PMID 29426988
25. Dai Y, Covarrubias D, Uppot R, et al. Image-Guided Percutaneous Radiofrequency Ablation of Central Renal Cell Carcinoma: Assessment of Clinical Efficacy and Safety in 31 Tumors. *J Vasc Interv Radiol*. Dec 2017; 28(12): 1643-1650. PMID 28673657
26. Dvorak P, Hoffmann P, Brodak M, et al. Percutaneous radiofrequency and microwave ablation in the treatment of renal tumors - 10 years of experience. *Wideochir Inne Tech Maloinwazyjne*. Dec 2017; 12(4): 394-402. PMID 29362655
27. Pantelidou M, Challacombe B, McGrath A, et al. Percutaneous Radiofrequency Ablation Versus Robotic-Assisted Partial Nephrectomy for the Treatment of Small Renal Cell Carcinoma. *Cardiovasc Intervent Radiol*. Nov 2016; 39(11): 1595-1603. PMID 27435582
28. Iannuccilli JD, Dupuy DE, Beland MD, et al. Effectiveness and safety of computed tomography-guided radiofrequency ablation of renal cancer: a 14-year single institution experience in 203 patients. *Eur Radiol*. Jun 2016; 26(6): 1656-64. PMID 26373755
29. Schlijper RC, Grutters JP, Houben R, et al. What to choose as radical local treatment for lung metastases from colo-rectal cancer: surgery or radiofrequency ablation?. *Cancer Treat Rev*. Feb 2014; 40(1): 60-7. PMID 23768754

30. Ratko TA, Vats V, Brock J, et al. Local Nonsurgical Therapies for Stage I and Symptomatic Obstructive Non- Small-Cell Lung Cancer (Comparative Effectiveness Review No. 112). Rockville, MD: Agency for Healthcare Research and Quality; 2013.
31. Bilal H, Mahmood S, Rajashanker B, et al. Is radiofrequency ablation more effective than stereotactic ablative radiotherapy in patients with early stage medically inoperable non-small cell lung cancer?. *Interact Cardiovasc Thorac Surg*. Aug 2012; 15(2): 258-65. PMID 22581864
32. Chan VO, McDermott S, Malone DE, et al. Percutaneous radiofrequency ablation of lung tumors: evaluation of the literature using evidence-based techniques. *J Thorac Imaging*. Feb 2011; 26(1): 18-26. PMID 20829720
33. Hasegawa T, Takaki H, Kodama H, et al. Three-year Survival Rate after Radiofrequency Ablation for Surgically Resectable Colorectal Lung Metastases: A Prospective Multicenter Study. *Radiology*. Mar 2020; 294(3): 686-695. PMID 31934829
34. Huang L, Han Y, Zhao J, et al. Is radiofrequency thermal ablation a safe and effective procedure in the treatment of pulmonary malignancies?. *Eur J Cardiothorac Surg*. Mar 2011; 39(3): 348-51. PMID 20663679
35. Zemlyak A, Moore WH, Bilfinger TV. Comparison of survival after sublobar resections and ablative therapies for stage I non-small cell lung cancer. *J Am Coll Surg*. Jul 2010; 211(1): 68-72. PMID 20610251
36. Lencioni R, Crocetti L, Cioni R, et al. Response to radiofrequency ablation of pulmonary tumours: a prospective, intention-to-treat, multicentre clinical trial (the RAPTURE study). *Lancet Oncol*. Jul 2008; 9(7): 621-8. PMID 18565793
37. Zhu JC, Yan TD, Glenn D, et al. Radiofrequency ablation of lung tumors: feasibility and safety. *Ann Thorac Surg*. Apr 2009; 87(4): 1023-8. PMID 19324122
38. Pennathur A, Abbas G, Gooding WE, et al. Image-guided radiofrequency ablation of lung neoplasm in 100 consecutive patients by a thoracic surgical service. *Ann Thorac Surg*. Nov 2009; 88(5): 1601-6; discussion 1607-8. PMID 19853119
39. Xia LY, Hu QL, Xu WY. Efficacy and Safety of Radiofrequency Ablation for Breast Cancer Smaller Than 2 cm: A Systematic Review and Meta-Analysis. *Front Oncol*. 2021; 11: 651646. PMID 34012918
40. Peek MCL, Ahmed M, Napoli A, et al. Minimally invasive ablative techniques in the treatment of breast cancer: a systematic review and meta-analysis. *Int J Hyperthermia*. Mar 2017; 33(2): 191-202. PMID 27575566
41. Zhao Z, Wu F. Minimally-invasive thermal ablation of early-stage breast cancer: a systemic review. *Eur J Surg Oncol*. Dec 2010; 36(12): 1149-55. PMID 20889281
42. Soukup B, Bismohun S, Reefy S, et al. The evolving role of radiofrequency ablation therapy of breast lesions. *Anticancer Res*. Sep 2010; 30(9): 3693-7. PMID 20944155
43. Ito T, Oura S, Nagamine S, et al. Radiofrequency Ablation of Breast Cancer: A Retrospective Study. *Clin Breast Cancer*. Aug 2018; 18(4): e495-e500. PMID 29079443
44. Li P, Xiao-Yin T, Cui D, et al. Evaluation of the safety and efficacy of percutaneous radiofrequency ablation for treating multiple breast fibroadenoma. *J Cancer Res Ther*. Dec 2016; 12(Supplement): C138-C142. PMID 28230006
45. Wilson M, Korourian S, Boneti C, et al. Long-term results of excision followed by radiofrequency ablation as the sole means of local therapy for breast cancer. *Ann Surg Oncol*. Oct 2012; 19(10): 3192-8. PMID 22911363
46. Kinoshita T, Iwamoto E, Tsuda H, et al. Radiofrequency ablation as local therapy for early breast carcinomas. *Breast Cancer*. Jan 2011; 18(1): 10-7. PMID 20072824
47. Imoto S, Wada N, Sakemura N, et al. Feasibility study on radiofrequency ablation followed by partial mastectomy for stage I breast cancer patients. *Breast*. Apr 2009; 18(2): 130-4. PMID 19324550
48. Garbay JR, Mathieu MC, Lamuraglia M, et al. Radiofrequency thermal ablation of breast cancer local recurrence: a phase II clinical trial. *Ann Surg Oncol*. Nov 2008; 15(11): 3222-6. PMID 18709415
49. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid*. Jan 2016; 26(1): 1-133. PMID 26462967
50. Xu X, Peng Y, Han G. Five-year follow-up results of thermal ablation for benign thyroid nodules: Systematic review and meta-analysis. *Am J Otolaryngol*. 2024; 45(1): 104025. PMID 37639985
51. Cho SJ, Baek JH, Chung SR, et al. Long-Term Results of Thermal Ablation of Benign Thyroid Nodules: A Systematic Review and Meta-Analysis. *Endocrinol Metab (Seoul)*. Jun 2020; 35(2): 339-350. PMID 32615718
52. Chen F, Tian G, Kong D, et al. Radiofrequency ablation for treatment of benign thyroid nodules: A PRISMA-compliant systematic review and meta-analysis of outcomes. *Medicine (Baltimore)*. Aug 2016; 95(34): e4659. PMID 27559968
53. Fuller CW, Nguyen SA, Lohia S, et al. Radiofrequency ablation for treatment of benign thyroid nodules: systematic review. *Laryngoscope*. Jan 2014; 124(1): 346-53. PMID 24122763
54. Kim JH, Yoo WS, Park YJ, et al. Efficacy and Safety of Radiofrequency Ablation for Treatment of Locally Recurrent Thyroid Cancers Smaller than 2 cm. *Radiology*. Sep 2015; 276(3): 909-18. PMID 25848897
55. Owen RP, Khan SA, Negassa A, et al. Radiofrequency ablation of advanced head and neck cancer. *Arch Otolaryngol Head Neck Surg*. May 2011; 137(5): 493-8. PMID 21576561
56. Brook AL, Gold MM, Miller TS, et al. CT-guided radiofrequency ablation in the palliative treatment of recurrent advanced head and neck malignancies. *J Vasc Interv Radiol*. May 2008; 19(5): 725-35. PMID 18440462
57. Owen RP, Silver CE, Ravikumar TS, et al. Techniques for radiofrequency ablation of head and neck tumors. *Arch Otolaryngol Head Neck Surg*. Jan 2004; 130(1): 52-6. PMID 14732768
58. Rey VE, Labrador R, Falcon M, et al. Transvaginal Radiofrequency Ablation of Myomas: Technique, Outcomes, and Complications. *J Laparoendosc Adv Surg Tech A*. Jan 2019; 29(1): 24-28. PMID 30198831
59. Yin G, Chen M, Yang S, et al. Treatment of uterine myomas by radiofrequency thermal ablation: a 10-year retrospective cohort study. *Reprod Sci*. May 2015; 22(5): 609-14. PMID 25355802

60. Liu B, Mo C, Wang W, et al. Treatment outcomes of percutaneous radiofrequency ablation versus adrenalectomy for adrenal metastases: a retrospective comparative study. *J Endocrinol Invest*. Sep 2020; 43(9): 1249-1257. PMID 32166699
61. Yang MH, Tyan YS, Huang YH, et al. Comparison of radiofrequency ablation versus laparoscopic adrenalectomy for benign aldosterone-producing adenoma. *Radiol Med*. Oct 2016; 121(10): 811-9. PMID 27300650
62. Locklin JK, Mannes A, Berger A, et al. Palliation of soft tissue cancer pain with radiofrequency ablation. *J Support Oncol*. 2004; 2(5): 439-45. PMID 15524075
63. Rosenthal DI. Radiofrequency treatment. *Orthop Clin North Am*. Jul 2006; 37(3): 475-84, viii. PMID 16846772
64. Liapi E, Geschwind JF. Transcatheter and ablative therapeutic approaches for solid malignancies. *J Clin Oncol*. Mar 10 2007; 25(8): 978-86. PMID 17350947
65. Spiliotis JD, Datsis AC, Michalopoulos NV, et al. Radiofrequency ablation combined with palliative surgery may prolong survival of patients with advanced cancer of the pancreas. *Langenbecks Arch Surg*. Jan 2007; 392(1): 55-60. PMID 17089173
66. Zou YP, Li WM, Zheng F, et al. Intraoperative radiofrequency ablation combined with 125 iodine seed implantation for unresectable pancreatic cancer. *World J Gastroenterol*. Oct 28 2010; 16(40): 5104-10. PMID 20976848
67. Cantore M, Girelli R, Mambrini A, et al. Combined modality treatment for patients with locally advanced pancreatic adenocarcinoma. *Br J Surg*. Aug 2012; 99(8): 1083-8. PMID 22648697
68. Rombouts SJ, Vogel JA, van Santvoort HC, et al. Systematic review of innovative ablative therapies for the treatment of locally advanced pancreatic cancer. *Br J Surg*. Feb 2015; 102(3): 182-93. PMID 25524417
69. Kameyama S, Murakami H, Masuda H, et al. Minimally invasive magnetic resonance imaging-guided stereotactic radiofrequency thermocoagulation for epileptogenic hypothalamic hamartomas. *Neurosurgery*. Sep 2009; 65(3): 438-49; discussion 449. PMID 19687687
70. Vavra P, Dostalík J, Zacharoulis D, et al. Endoscopic radiofrequency ablation in colorectal cancer: initial clinical results of a new bipolar radiofrequency ablation device. *Dis Colon Rectum*. Feb 2009; 52(2): 355-8. PMID 19279436
71. Mylona S, Karagiannis G, Patsoura S, et al. Palliative treatment of rectal carcinoma recurrence using radiofrequency ablation. *Cardiovasc Intervent Radiol*. Aug 2012; 35(4): 875-82. PMID 22167304
72. Ripley RT, Gajdos C, Reppert AE, et al. Sequential radiofrequency ablation and surgical debulking for unresectable colorectal carcinoma: thermo-surgical ablation. *J Surg Oncol*. Feb 2013; 107(2): 144-7. PMID 22927225
73. Howington JA, Blum MG, Chang AC, et al. Treatment of stage I and II non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. May 2013; 143(5 Suppl): e278S-e313S. PMID 23649443
74. Donington J, Ferguson M, Mazzone P, et al. American College of Chest Physicians and Society of Thoracic Surgeons consensus statement for evaluation and management for high-risk patients with stage I non-small cell lung cancer. *Chest*. Dec 2012; 142(6): 1620-1635. PMID 23208335
75. Orloff LA, Noel JE, Stack BC, et al. Radiofrequency ablation and related ultrasound-guided ablation technologies for treatment of benign and malignant thyroid disease: An international multidisciplinary consensus statement of the American Head and Neck Society Endocrine Surgery Section with the Asia Pacific Society of Thyroid Surgery, Associazione Medici Endocrinologi, British Association of Endocrine and Thyroid Surgeons, European Thyroid Association, Italian Society of Endocrine Surgery Units, Korean Society of Thyroid Radiology, Latin American Thyroid Society, and Thyroid Nodules Therapies Association. *Head Neck*. Mar 2022; 44(3): 633-660. PMID 34939714
76. Campbell S, Uzzo RG, Allaf ME, et al. Renal Mass and Localized Renal Cancer: AUA Guideline. *J Urol*. Sep 2017; 198(3): 520-529. PMID 28479239
77. Campbell SC, Clark PE, Chang SS, et al. Renal Mass and Localized Renal Cancer: Evaluation, Management, and Follow-Up: AUA Guideline: Part I. *J Urol*. Aug 2021; 206(2): 199-208. PMID 34115547
78. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Non-small cell lung cancer. Version 7.2024. Updated June 26, 2024. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed July 25, 2024.
79. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Thyroid Carcinoma. Version 3.2024. Updated June 18, 2024. https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf. Accessed July 26, 2024.
80. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Kidney Cancer. Version 1.2025. Updated July 1, 2024. https://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf. Accessed July 24, 2024.
81. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Colon Cancer. Version 4.2024. Updated July 3, 2024. https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed July 23, 2024.
82. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Head and Neck Cancers. Version 4.2024. Updated May 1, 2024. https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf. Accessed July 27, 2024.
83. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 4.2024. Updated July 3, 2024. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed July 22, 2024.
84. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Bone Cancer. Version 2.2024. Updated March 12, 2024. https://www.nccn.org/professionals/physician_gls/pdf/bone.pdf. Accessed July 21, 2024.
85. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Pancreatic Adenocarcinoma. Version 2.2024. Updated April 30, 2024. https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf. Accessed July 28, 2024.
86. National Institute for Health and Care Excellence (NICE). Computed tomography-guided thermocoagulation of osteoid osteoma [IPG53]. 2004; <https://www.nice.org.uk/guidance/ipg53>. Accessed July 30, 2024.
87. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation of renal cancer [IPG353]. 2010; <https://www.nice.org.uk/guidance/ipg353>. Accessed July 28, 2024.
88. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation for primary and secondary lung cancers [IPG372]. 2010; <https://www.nice.org.uk/guidance/ipg372>. Accessed July 29, 2024.

89. National Institute for Health and Care Excellence (NICE). Ultrasound-guided percutaneous radiofrequency ablation for benign thyroid nodules [IPG562]. 2016; <https://www.nice.org.uk/guidance/IPG562>. Accessed July 27, 2024.
90. Morris CS, Baerlocher MO, Dariushnia SR, et al. Society of Interventional Radiology Position Statement on the Role of Percutaneous Ablation in Renal Cell Carcinoma: Endorsed by the Canadian Association for Interventional Radiology and the Society of Interventional Oncology. *J Vasc Interv Radiol*. Feb 2020; 31(2): 189-194.e3. PMID 31917025

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

| Date | Action | Description |
|----------------|----------------|---|
| June 2012 | New policy | Medically necessary with criteria for: osteolytic bone metastases, osteoid osteomas, localized renal cell carcinoma, isolated peripheral NSCLC, malignant nonpulmonary tumors metastatic to lung; investigational for: breast tumors, lung, renal, osteoid osteomas, and bony metastases not meeting criteria and all other tumors. |
| March 2013 | Replace policy | Policy updated with literature review, reworded not medically necessary policy statement and added thyroid as not medically necessary. References 12, 19, 29, 42, 50-52, 59, 62-63 added. Other references deleted. |
| March 2014 | Replace policy | Policy updated with literature review, Policy statements unchanged. References added, renumbered or removed. |
| March 2015 | Replace policy | Policy updated with literature review; references 4, 48, and 56 added; other references deleted. Policy statements unchanged |
| December 2016 | Replace policy | Policy updated with literature review; reference 13 added; references 54- 56 updated; some references removed. Policy statements unchanged. |
| December 2017 | Replace policy | Policy updated with literature review through July 20, 2017; reference 59 added. Policy statements unchanged. |
| December 2018 | Replace policy | Policy updated with literature review through July 26, 2018; references 5- 7, 15-19, 31, 34-35, 40, 42, and 50 added. Policy statements unchanged. |
| September 2019 | Replace policy | Policy updated with literature review through July 8, 2019; references added, references on NCCN updated. Policy statements unchanged. |
| December 2020 | Replace policy | Policy updated with literature review through July 27, 2020; references added. Policy statements unchanged. |
| December 2021 | Replace policy | Policy updated with literature review through July 28, 2021; references added. Policy statements unchanged. |
| December 2022 | Replace policy | Policy updated with literature review through August 10, 2022; references added. Minor editorial refinements to policy statements; intent unchanged. |
| December 2023 | Replace policy | Policy updated with literature review through August 7, 2023; no references added. Policy statements unchanged. |
| December 2024 | Replace policy | Policy updated with literature review through July 31, 2024; reference added. Policy statements unchanged. |

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.