

Federal Employee Program.
Blue Cross Blue Shield Association 750 9th St NW, Suite 900
Washington, D.C. 20001
1-800-624-5060
Fax 1-877-378-4727

5.21.162

Section: Prescription Drugs Effective Date: January 1, 2025

Subsection: Antineoplastic Agents Original Policy Date: September 25, 2020

Subject: Gavreto Page: 1 of 5

Last Review Date: December 13, 2024

Gavreto

Description

Gavreto (pralsetinib)

Background

Gavreto (pralsetinib) is a kinase inhibitor of wild-type *RET* and oncogenic *RET* fusions and mutations. Certain *RET* fusion proteins and activating point mutations can drive tumorigenic potential through hyperactivation of downstream signaling pathways leading to uncontrolled cell proliferation. Gavreto exhibits anti-tumor activity in models harboring oncogenic *RET* fusions or mutations (1).

Regulatory Status

FDA-approved indications: Gavreto is a kinase inhibitor indicated for the treatment of: (1)

- Adult patients with metastatic rearranged during transfection (RET) fusion-positive nonsmall cell lung cancer (NSCLC) as detected by an FDA approved test
- Adult and pediatric patients 12 years of age and older with advanced or metastatic RET fusion-positive thyroid cancer who require systemic therapy and who are radioactive iodine-refractory (if refractory iodine is appropriate)

Patients should be selected for treatment with Gavreto based on the presence of a *RET* gene fusion (NSCLC or thyroid cancer) (1).

Gavreto has warnings regarding hepatotoxicity and hypertension. AST and ALT should be monitored prior to initiating Gavreto, every 2 weeks during the first 3 months, then monthly thereafter and as clinically indicated. Gavreto should not be initiated in patients with

5.21.162

Section: Prescription Drugs Effective Date: January 1, 2025

Subsection: Antineoplastic Agents Original Policy Date: September 25, 2020

Subject: Gavreto Page: 2 of 5

uncontrolled hypertension and blood pressure should be optimized prior to initiation. Blood pressure should be monitored after 1 week, at least monthly thereafter and as clinically indicated (1).

Gavreto can cause fetal harm when administered to a pregnant woman. Females of reproductive potential should be advised to use effective non-hormonal contraception during treatment with Gavreto and for 2 weeks after the final dose. Males with female partners of reproductive potential should be advised to use effective contraception during treatment with Gavreto and for 1 week after the final dose (1).

The safety and effectiveness of Gavreto have not been established in pediatric patients less than 18 years of age with *RET* fusion-positive NSCLC or in pediatric patients less than 12 years of age with *RET* fusion-positive thyroid cancer (1).

Related policies

Retevmo

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Gavreto may be considered **medically necessary** if the conditions indicated below are met.

Gavreto may be considered **investigational** for all other indications.

Prior-Approval Requirements

Diagnoses

Patient must have **ONE** of the following:

- 1. Metastatic non-small cell lung cancer (NSCLC)
 - a. 18 years of age or older
 - b. RET fusion-positive detected by an FDA approved test
- 2. Advanced or metastatic thyroid cancer
 - a. 12 years of age or older
 - b. *RET* fusion-positive and patient requires systemic therapy

5.21.162

Section: Prescription Drugs Effective Date: January 1, 2025

Subsection: Antineoplastic Agents Original Policy Date: September 25, 2020

Subject: Gavreto Page: 3 of 5

c. Radioactive iodine-refractory (if radioactive iodine is appropriate)

AND ALL of the following:

- a. Prescriber agrees to monitor AST, ALT, and blood pressure
- Females of reproductive potential only: patient will be advised to use effective non-hormonal contraception during treatment with Gavreto and for 2 weeks after the final dose
- Males with female partners of reproductive potential only: patient will be advised to use effective contraception during treatment with Gavreto and for 1 week after the final dose

Prior - Approval Renewal Requirements

Diagnoses

Patient must have **ONE** of the following:

- 1. Metastatic non-small cell lung cancer (NSCLC)
 - a. 18 years of age or older
- 2. Advanced or metastatic thyroid cancer
 - a. 12 years of age or older

AND ALL of the following:

- a. NO disease progression or unacceptable toxicity
- b. Prescriber agrees to monitor AST, ALT, and blood pressure
- Females of reproductive potential only: patient will be advised to use
 effective non-hormonal contraception during treatment with Gavreto and for 2
 weeks after the final dose
- Males with female partners of reproductive potential only: patient will be advised to use effective contraception during treatment with Gavreto and for 1 week after the final dose

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

Section: Prescription Drugs Effective Date: January 1, 2025

Subsection: Antineoplastic Agents Original Policy Date: September 25, 2020

Subject: Gavreto Page: 4 of 5

Quantity 360 capsules per 90 days

Duration 12 months

Prior - Approval Renewal Limits

Same as above

Rationale

Summary

Gavreto (pralsetinib) is a kinase inhibitor of wild-type *RET* and oncogenic *RET* fusions and mutations. Certain *RET* fusion proteins and activating point mutations can drive tumorigenic potential through hyperactivation of downstream signaling pathways leading to uncontrolled cell proliferation. Gavreto exhibits anti-tumor activity in models harboring oncogenic *RET* fusions or mutations (1).

Prior authorization is required to ensure the safe, clinically appropriate, and cost-effective use of Gavreto while maintaining optimal therapeutic outcomes.

References

- 1. Gavreto [package insert]. South San Fracisco, CA: Rigel Pharmaceuticals, Inc.; June 2024.
- 2. NCCN Drugs & Biologics Compendium[®] Pralsetinib 2024. National Comprehensive Cancer Network, Inc. Accessed on October 29, 2024.

Policy History	
Date	Action
September 2020 December 2020 January 2021 March 2021 March 2022 June 2023 September 2023 December 2023 March 2024 December 2024	Addition to PA Annual review Addition of indications: medullary thyroid cancer and thyroid cancer Annual review Annual review and reference update Annual review and reference update Per PI update, removed indication for MTC Annual review and reference update

5.21.162

Section: Prescription Drugs Effective Date: January 1, 2025

Subsection: Antineoplastic Agents Original Policy Date: September 25, 2020

Subject: Gavreto Page: 5 of 5

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on December 13, 2024 and is effective on January 1, 2025.