



FEP Medical Policy Manual

FEP 7.01.58 Intraoperative Neurophysiologic Monitoring

Effective Policy Date: July 1, 2023

Original Policy Date: December 2011

Related Policies:

None

Intraoperative Neurophysiologic Monitoring

Description

Description

Intraoperative neurophysiologic monitoring describes a variety of procedures used to monitor the integrity of neural pathways during high-risk neurosurgical, orthopedic, and vascular surgeries. It involves the detection of electrical signals produced by the nervous system in response to sensory or electrical stimuli to provide information about the functional integrity of neuronal structures. This evidence review does not address established neurophysiologic monitoring (ie, somatosensory-evoked potentials, motor-evoked potentials using transcranial electrical stimulation, brainstem auditory-evoked potentials, electromyography of cranial nerves, electroencephalography, electrocorticography), during spinal, intracranial, or vascular procedures.

OBJECTIVE

The objective of this evidence review is to determine whether neurophysiologic monitoring improves the net health outcome in individuals during surgeries that could damage their recurrent laryngeal nerve or peripheral nerves.

POLICY STATEMENT

Intraoperative neurophysiologic monitoring, which includes somatosensory-evoked potentials, motor-evoked potentials using transcranial electrical stimulation, brainstem auditory-evoked potentials, electromyography of cranial nerves, electroencephalography, and electrocorticography, may be considered **medically necessary** during spinal, intracranial, or vascular procedures.

Intraoperative neurophysiologic monitoring of the recurrent laryngeal nerve may be considered **medically necessary** in individuals undergoing:

- high-risk thyroid or parathyroid surgery, including:
 - total thyroidectomy
 - repeat thyroid or parathyroid surgery
 - surgery for cancer
 - thyrotoxicosis
 - retrosternal or giant goiter
 - thyroiditis.
- anterior cervical spine surgery associated with any of the following increased risk situations:
 - prior anterior cervical surgery, particularly revision anterior cervical discectomy and fusion, revision surgery through a scarred surgical field, reoperation for pseudarthrosis, or revision for failed fusion
 - multilevel anterior cervical discectomy and fusion
 - preexisting recurrent laryngeal nerve pathology, when there is residual function of the recurrent laryngeal nerve.

Intraoperative neurophysiologic monitoring of the recurrent laryngeal nerve during anterior cervical spine surgery not meeting the criteria above or during esophageal surgeries is considered **investigational**.

Intraoperative monitoring of visual-evoked potentials is considered **investigational**.

Due to the lack of monitors approved by the U.S. Food and Drug Administration, intraoperative monitoring of motor-evoked potentials using transcranial magnetic stimulation is considered **investigational**.

Intraoperative electromyography and nerve conduction velocity monitoring during surgery on the peripheral nerves is considered **not medically necessary**.

Note: These policy statements refer only to use of these techniques as part of intraoperative monitoring. Other clinical applications of these techniques, such as visual-evoked potentials and electromyography, are not considered in this policy.

POLICY GUIDELINES

Intraoperative neurophysiologic monitoring, including somatosensory-evoked potentials and motor-evoked potentials using transcranial electrical stimulation, brainstem auditory-evoked potentials, electromyography of cranial nerves, electroencephalography, and electrocorticography, has broad acceptance, particularly for spine surgery and open abdominal aorta aneurysm repairs. Therefore, this evidence review focuses on monitoring of the recurrent laryngeal nerve during neck surgeries and monitoring of peripheral nerves.

Constant communication among the surgeon, neurophysiologist, and anesthetist is required for safe and effective intraoperative neurophysiologic monitoring.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

Intraoperative neurophysiologic monitoring is considered reimbursable as a separate service only when a licensed health care practitioner, other than the operating surgeon, interprets the monitoring. The monitoring is performed by a health care practitioner or technician who is in attendance in the operating room throughout the procedure.

Implementation of a local policy on this technology may also involve discussions about credentialing of those providing the intraoperative monitoring services, as well as on-site versus remote real-time review and interpretation.

Coding for intraoperative monitoring uses time-based codes; they are not based on the number (single vs. multiple) of modalities used.

FDA REGULATORY STATUS

A number of EEG and EMG monitors have been cleared for marketing by the FDA through the 510(k) process.

FDA product code: GWQ.

Intraoperative neurophysiologic monitoring of motor-evoked potentials using transcranial magnetic stimulation does not have FDA approval.

RATIONALE

Summary of Evidence

For individuals who are undergoing thyroid or parathyroid surgery and are at high risk of injury to the recurrent laryngeal nerve who receive intraoperative neurophysiologic monitoring, the evidence includes a large randomized controlled trial (RCT) and systematic reviews. Relevant outcomes are morbid events, functional outcomes, and quality of life. The strongest evidence on neurophysiologic monitoring derives from a RCT of 1000 patients undergoing thyroid surgery. This RCT found a significant reduction in recurrent laryngeal nerve injury in patients at high-risk for injury. High-risk in this trial was defined as surgery for cancer, thyrotoxicosis, retrosternal or giant goiter, or thyroiditis. The high-risk category may also include patients with prior thyroid or parathyroid surgery or total thyroidectomy. A low volume of surgeries might also contribute to a higher risk for recurrent laryngeal nerve injury. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing anterior cervical spine surgery and are at high-risk of injury to the recurrent laryngeal nerve who receive intraoperative neurophysiologic monitoring, the evidence includes 3 systematic reviews of case series and cohort studies. Relevant outcomes are morbid events, functional outcomes, and quality of life. Two of the 3 analyses compared the risk of nerve injury using intraoperative neurophysiologic monitoring with no intraoperative neurophysiologic monitoring and found no statistically significant difference. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing esophageal surgery who receive intraoperative neurophysiologic monitoring, the evidence includes a systematic review of mainly nonrandomized comparative studies. Relevant outcomes are morbid events, functional outcomes, and quality of life. The systematic review found less recurrent laryngeal nerve palsy with intraoperative neurophysiologic monitoring but conclusions are limited by the design of the included studies. Current evidence is not sufficiently robust to determine whether neurophysiologic monitoring reduces recurrent laryngeal nerve injury in patients undergoing esophageal surgery. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing surgery proximal to a peripheral nerve who receive intraoperative neurophysiologic monitoring, the evidence includes case series and a controlled cohort study. Relevant outcomes are morbid events, functional outcomes, and quality of life. Surgical guidance with peripheral intraoperative neurophysiologic monitoring and the predictive ability of monitoring of peripheral nerves have been reported. No prospective comparative studies were identified that assessed whether outcomes are improved with neurophysiologic monitoring. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are undergoing spinal instrumentation requiring screws or distraction who receive intraoperative neurophysiologic monitoring, the evidence includes systematic reviews of nonrandomized studies. Relevant outcomes are morbid events, functional outcomes, and quality of life. The available evidence suggests that intraoperative neurophysiologic monitoring has high sensitivity and specificity for detecting neurologic deficits. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in "Supplemental Information" if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Neurology

In 1990 (updated in 2012), the American Academy of Neurology (AAN) published an assessment of intraoperative neurophysiologic monitoring, with an evidence-based guideline update by the AAN and the American Clinical Neurophysiology Society (ACNS) in 2012 (guideline last reaffirmed on October 17, 2020).^{1,2} The 1990 assessment indicated that monitoring requires a team approach with a well-trained physician-neurophysiologist to provide or supervise monitoring. Electroencephalogram (EEG) monitoring is used during carotid endarterectomy or for other similar situations in which cerebral blood flow is at high risk. Electrocorticography from surgically exposed cortex can help to define the optimal limits of surgical resection or identify regions of greatest impairment, while sensory cortex somatosensory-evoked potentials can help to localize the central fissure and motor cortex. Auditory-evoked potentials, along with cranial nerve monitoring can be used during posterior fossa neurosurgical procedures. Spinal cord somatosensory-evoked potentials are frequently used to monitor the spinal cord during orthopedic or neurosurgical procedures around the spinal cord, or cross-clamping of the thoracic aorta. Electromyographic monitoring during procedures near the roots and peripheral nerves can be used to warn of excessive traction or other impairment of motor nerves. At the time of the 1990 assessment, motor-evoked potentials were considered investigational by many neurophysiologists. The 2012 update, which was endorsed by the American Association of Neuromuscular & Electrodiagnostic Medicine (AANEM), concluded that the available evidence supported intraoperative neurophysiologic monitoring using somatosensory-evoked potentials or motor-evoked potentials when conducted under the supervision of a clinical neurophysiologist experienced with intraoperative neurophysiologic monitoring. Evidence was insufficient to evaluate intraoperative neurophysiologic monitoring when conducted by technicians alone or by an automated device.

In 2012, the AAN published a model policy on principles of coding for intraoperative neurophysiologic monitoring and testing (last amended July 31, 2018).⁶³ The background section of this document provides the following information on the value of intraoperative neurophysiologic monitoring in averting neural injuries during surgery:

1. "Value of EEG Monitoring in Carotid Surgery. Carotid occlusion, incident to carotid endarterectomies, poses a high-risk for cerebral hemispheric injury. Electroencephalogram (EEG) monitoring is capable of detecting cerebral ischemia, a serious prelude to injury. Studies of continuous monitoring established the ability of electroencephalogram EEG to correctly predict risks of postoperative deficits after a deliberate, but necessary, carotid occlusion as part of the surgical procedure. The surgeon can respond to adverse EEG events by raising blood pressure, implanting a shunt, adjusting a poorly functioning shunt, or performing other interventions.
2. Multicenter Data in Spinal Surgeries. An extensive multicenter study conducted in 1995 demonstrated that [intraoperative neurophysiologic monitoring] using [sensory-evoked potentials] reduced the risk of paraplegia by 60% in spinal surgeries. The incidence of false negative cases, wherein an operative complication occurred without having been detected by the monitoring procedure, was small: 0.06%.
3. Technology Assessment of Monitoring in Spinal Surgeries. A technology assessment by the McGill University Health Center...reviewed 11 studies and concluded that spinal [intraoperative neurophysiologic monitoring] is capable of substantially reducing injury in surgeries that pose a risk to spinal cord integrity. It recommended combined sensory-evoked potentials/motor-evoked potential monitoring, under the presence or constant availability of a monitoring physician, for all cases of spinal surgery for which there is a risk of spinal cord injury.
4. Value of Combined Motor and Sensory Monitoring. Numerous studies of post-surgical paraparesis and quadriparesis have shown that both sensory-evoked potentials and motor-evoked potential monitoring had predicted adverse outcomes in a timely fashion. The timing of the predictions allowed the surgeons the opportunity to intervene and prevent adverse outcomes. The 2 different techniques (sensory-evoked potentials and motor-evoked potential) monitor different spinal cord tracts. Sometimes, one of the techniques cannot be used for practical purposes, for anesthetic reasons, or because of preoperative absence of signals in those pathways. Thus, the decision about which of these techniques to use needs to be tailored to the individual patient's circumstances.
5. Protecting the Spinal Cord from Ischemia during Aortic Procedures. Studies have shown that [intraoperative neurophysiologic monitoring] accurately predicts risks for spinal cord ischemia associated with clamping the aorta or ligating segmental spinal arteries. [Intraoperative neurophysiologic monitoring] can assess whether the spinal cord is tolerating the degree of relative ischemia in these procedures. The surgeon can then respond by raising blood pressure, implanting a shunt, re-implanting segmental vessels, draining spinal fluid, or through other interventions...

6. Value of EMG [electromyogram] monitoring. Selective posterior rhizotomy in cerebral palsy significantly reduces spasticity, increases range of motion, and improves functional skills. Electromyography during this procedure can assist in selecting specific dorsal roots to transect. Electromyogram (EMG) can also be used in peripheral nerve procedures that pose a risk of injuries to nerves...

7. Value of Spinal Monitoring using somatosensory-evoked potentials and motor-evoked potentials. According to a recent review of spinal monitoring using somatosensory-evoked potential and motor-evoked potentials by the Therapeutics and Technology Assessment Subcommittee of AAN and ACNS, [intraoperative neurophysiologic monitoring] is established as effective to predict an increased risk of the adverse outcomes of paraparesis, paraplegia, and quadriplegia in spinal surgery (4 Class I and 7 Class II studies). Surgeons and other members of the operating team should be alerted to the increased risk of severe adverse neurologic outcomes in patients with important [intraoperative neurophysiologic monitoring] changes (Level A)."

The AAN model policy also offered guidance on personnel and monitoring standards for intraoperative neurophysiologic monitoring and somatosensory-evoked potential.

American Association of Neurological Surgeons and Congress of Neurological Surgeons

In 2018, the American Association of Neurological Surgeons (AANS) and Congress of Neurological Surgeons updated their position statement on intraoperative neurophysiologic monitoring during routine spinal surgery.⁶⁴ They stated that intraoperative neurophysiologic monitoring, especially motor evoked potential, "is a reliable diagnostic tool for assessment of spinal cord integrity during surgery" (Level 1 evidence). Intraoperative motor evoked potentials may also "predict recovery in traumatic cervical spinal cord injury." However, AANS and Congress of Neurological Surgeons found no evidence that such monitoring provides a therapeutic benefit. The statement also recommends that intraoperative neurophysiologic monitoring should be used when the operating surgeon believes it is warranted for diagnostic value, such as with "deformity correction, spinal instability, spinal cord compression, intradural spinal cord lesions, and when in proximity to peripheral nerves or roots." In addition, they recommend spontaneous and evoked electromyography "for minimally invasive lateral retroperitoneal transpsoas approaches to the lumbar spine" and during screw insertion.

In 2014, the same organizations published guidance on electrophysiological monitoring for lumbar fusion procedures.⁶⁵ The authors concluded that there was a lack of high quality studies and that routine intraoperative monitoring during lumbar fusion could not be recommended. Evidence regarding the efficacy of intraoperative monitoring to recover nerve function or affect the outcome of surgery.

American Association of Neuromuscular & Electrodiagnostic Medicine

In 2023, the AANEM updated their position statement on electrodiagnostic medicine.⁵ The recommendations indicated that intraoperative sensory-evoked potentials have demonstrated usefulness for monitoring of spinal cord, brainstem, and brain sensory tracts. The AANEM stated that intraoperative somatosensory-evoked potential monitoring is indicated for select spine surgeries in which there is a risk of additional nerve root or spinal cord injury. Indications for somatosensory-evoked potential monitoring may include, but are not limited to, complex, extensive, or lengthy procedures, and when mandated by hospital policy. However, intraoperative somatosensory-evoked potential monitoring may not be indicated for routine lumbar or cervical root decompression.

American Clinical Neurophysiology Society

In 2009, the ACNS recommended standards for intraoperative neurophysiologic monitoring.⁴ **Guideline 11A included the following statement⁶⁶:**

"The monitoring team should be under the direct supervision of a physician with training and experience in neurophysiologic intraoperative monitoring. The monitoring physician should be licensed in the state and privileged to interpret neurophysiologic testing in the hospital in which the surgery is being performed. He/she is responsible for real-time interpretation of neurophysiologic intraoperative monitoring data. The monitoring physician should be present in the operating room or have access to intraoperative neurophysiologic monitoring data in real-time from a remote location and be in communication with the staff in the operating room. There are many methods of remote monitoring, however any method used must conform to local and national protected health information guidelines. The specifics of this availability (ie, types of surgeries) should be decided by the hospital credentialing committee. In order to devote the needed attention, it is recommended that the monitoring physician interpret no more than three cases concurrently."

American Head and Neck Society

In 2022, the American Head and Neck Society Endocrine Surgery Section and the International Neural Monitoring Study Group published a clinical review of intraoperative nerve monitoring during pediatric thyroid surgery.⁶⁷ The review stated that intraoperative neurophysiologic monitoring can be considered in all pediatric thyroid surgeries. Procedures for which monitoring may be most beneficial include: total thyroidectomy, hemithyroidectomy in which the contralateral vocal cord is paralyzed, and reoperative surgeries.

American Society of Neurophysiological Monitoring

In 2018, the American Society of Neurophysiological Monitoring (ASNM) published practice guidelines for the supervising professional on intraoperative neurophysiologic monitoring.¹⁵ The ASNM (2013) position statement on intraoperative motor-evoked potential monitoring indicated that motor-evoked potentials are an established practice option for cortical and subcortical mapping and monitoring during surgeries risking motor injury in the brain, brainstem, spinal cord, or facial nerve.⁶⁸

Scoliosis Research Society

In 2020, the Scoliosis Research Society published an information statement on neurophysiologic monitoring during spinal deformity surgery.⁶⁹ The Society concluded that neurophysiologic monitoring can allow for early detection of complications and possibly prevent postoperative neurologic injury, and is considered optimal care when the spinal cord is at risk, which warrants a strong recommendation unless there are contraindications. The standard method of intraoperative monitoring should include transcranial motor evoked potentials and somatosensory evoked potentials with or without electromyographic monitoring.

National Institute for Health and Care Excellence

In 2008, a guidance from NICE on intraoperative neurophysiologic monitoring during thyroid surgery found no major safety concerns.⁷⁰ Regarding efficacy, intraoperative neurophysiologic monitoring was indicated as helpful "in performing more complex operations such as reoperative surgery and operations on large thyroid glands."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

The Centers for Medicare & Medicaid Services has indicated that EEG monitoring "may be covered routinely in carotid endarterectomies and in other neurological procedures where cerebral perfusion could be reduced. Such other procedures might include aneurysm surgery where hypotensive anesthesia is used or other cerebral vascular procedures where cerebral blood flow may be interrupted."⁷¹ Coverage determinations for other modalities were not identified.

The Centers for Medicare & Medicaid Services Physician Fee Schedule Final Rule (2013) discussed payment of neurophysiologic monitoring. The rule states that CPT code 95940, which is reported when a physician monitors a patient directly, is payable by Medicare. CPT code 95941, which is used for remote monitoring, was made invalid for submission to Medicare.

In the Final Rule, the Centers established a HCPCS G code (see Policy Guidelines section) for reporting physician monitoring performed from outside of the operating room (nearby or remotely). HCPCS code G0453 "may be billed only for undivided attention by the monitoring physician to a single beneficiary [1:1 technologist to oversight physician billing], and not for simultaneous attention by the monitoring physician to more than one patient."⁷²

REFERENCES

1. Assessment: intraoperative neurophysiology. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. Nov 1990; 40(11): 1644-6. PMID 2234418
2. Nuwer MR, Emerson RG, Galloway G, et al. Evidence-based guideline update: intraoperative spinal monitoring with somatosensory and transcranial electrical motor evoked potentials: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology and the American Clinical Neurophysiology Society. *Neurology*. Feb 21 2012; 78(8): 585-9. PMID 22351796
3. Skinner SA, Cohen BA, Morledge DE, et al. Practice guidelines for the supervising professional: intraoperative neurophysiological monitoring. *J Clin Monit Comput*. Apr 2014; 28(2): 103-11. PMID 24022172
4. American Clinical Neurophysiology Society. ACNS Guidelines and Consensus Statements. <http://www.acns.org/practice/guidelines>. Accessed March 3, 2023.
5. American Association of Neuromuscular & Electrodiagnostic Medicine. Position Statement: Recommended Policy for Electrodiagnostic Medicine. updated 2023; <https://www.aanem.org/Advocacy/Position-Statements/Recommended-Policy-for-Electrodiagnostic-Medicine>. Accessed March 1, 2023.
6. Resnick DK, Choudhri TF, Dailey AT, et al. Guidelines for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 15: electrophysiological monitoring and lumbar fusion. *J Neurosurg Spine*. Jun 2005; 2(6): 725-32. PMID 16028743
7. Henry BM, Graves MJ, Vikse J, et al. The current state of intermittent intraoperative neural monitoring for prevention of recurrent laryngeal nerve injury during thyroidectomy: a PRISMA-compliant systematic review of overlapping meta-analyses. *Langenbecks Arch Surg*. Jun 2017; 402(4): 663-673. PMID 28378238
8. Pisanu A, Porceddu G, Podda M, et al. Systematic review with meta-analysis of studies comparing intraoperative neuromonitoring of recurrent laryngeal nerves versus visualization alone during thyroidectomy. *J Surg Res*. May 01 2014; 188(1): 152-61. PMID 24433869
9. Sun W, Liu J, Zhang H, et al. A meta-analysis of intraoperative neuromonitoring of recurrent laryngeal nerve palsy during thyroid reoperations. *Clin Endocrinol (Oxf)*. Nov 2017; 87(5): 572-580. PMID 28585717
10. Pardal-Refoyo JL, Ochoa-Sangrador C. Bilateral recurrent laryngeal nerve injury in total thyroidectomy with or without intraoperative neuromonitoring. Systematic review and meta-analysis. *Acta Otorrinolaringol Esp*. 2016; 67(2): 66-74. PMID 26025358
11. Barczyński M, Konturek A, Cichoń S. Randomized clinical trial of visualization versus neuromonitoring of recurrent laryngeal nerves during thyroidectomy. *Br J Surg*. Mar 2009; 96(3): 240-6. PMID 19177420
12. Vasileiadis I, Karatzas T, Charitoudis G, et al. Association of Intraoperative Neuromonitoring With Reduced Recurrent Laryngeal Nerve Injury in Patients Undergoing Total Thyroidectomy. *JAMA Otolaryngol Head Neck Surg*. Oct 01 2016; 142(10): 994-1001. PMID 27490310
13. Ajiboye RM, Zoller SD, Sharma A, et al. Intraoperative Neuromonitoring for Anterior Cervical Spine Surgery: What Is the Evidence?. *Spine (Phila Pa 1976)*. Mar 15 2017; 42(6): 385-393. PMID 27390917
14. Erwood MS, Hadley MN, Gordon AS, et al. Recurrent laryngeal nerve injury following reoperative anterior cervical discectomy and fusion: a meta-analysis. *J Neurosurg Spine*. Aug 2016; 25(2): 198-204. PMID 27015129
15. Daniel JW, Botelho RV, Milano JB, et al. Intraoperative Neurophysiological Monitoring in Spine Surgery: A Systematic Review and Meta-Analysis. *Spine (Phila Pa 1976)*. Aug 2018; 43(16): 1154-1160. PMID 30063222
16. Chen B, Yang T, Wang W, et al. Application of Intraoperative Neuromonitoring (IONM) of the Recurrent Laryngeal Nerve during Esophagectomy: A Systematic Review and Meta-Analysis. *J Clin Med*. Jan 10 2023; 12(2). PMID 36675495
17. Komatsu S, Konishi T, Matsubara D, et al. Continuous Recurrent Laryngeal Nerve Monitoring During Single-Port Mediastinoscopic Radical Esophagectomy for Esophageal Cancer. *J Gastrointest Surg*. Dec 2022; 26(12): 2444-2450. PMID 36221021
18. Huang CL, Chen CM, Hung WH, et al. Clinical Outcome of Intraoperative Recurrent Laryngeal Nerve Monitoring during Thoracoscopic Esophagectomy and Mediastinal Lymph Node Dissection for Esophageal Cancer. *J Clin Med*. Aug 23 2022; 11(17). PMID 36078880
19. Zhao L, He J, Qin Y, et al. Application of intraoperative nerve monitoring for recurrent laryngeal nerves in minimally invasive McKeown esophagectomy. *Dis Esophagus*. Jul 12 2022; 35(7). PMID 34864953
20. Yuda M, Nishikawa K, Ishikawa Y, et al. Intraoperative nerve monitoring during esophagectomy reduces the risk of recurrent laryngeal nerve palsy. *Surg Endosc*. Jun 2022; 36(6): 3957-3964. PMID 34494155
21. Takeda S, Iida M, Kanekiyo S, et al. Efficacy of intraoperative recurrent laryngeal neuromonitoring during surgery for esophageal cancer. *Ann Gastroenterol Surg*. Jan 2021; 5(1): 83-92. PMID 33532684
22. Fujimoto D, Taniguchi K, Kobayashi H. Intraoperative neuromonitoring during prone thoracoscopic esophagectomy for esophageal cancer reduces the incidence of recurrent laryngeal nerve palsy: a single-center study. *Updates Surg*. Apr 2021; 73(2): 587-595. PMID 33415692
23. Kobayashi H, Kondo M, Mizumoto M, et al. Technique and surgical outcomes of mesenterization and intra-operative neural monitoring to reduce recurrent laryngeal nerve paralysis after thoracoscopic esophagectomy: A cohort study. *Int J Surg*. Aug 2018; 56: 301-306. PMID 29879478
24. Zhu W, Yang F, Cao J, Zhao C, Dong B, Chen D. Application of recurrent laryngeal nerve detector in the neck anastomosis of upper or middle-thoracic esophageal carcinoma. *Cancer Res Clin*. 2018;30:233236.
25. Hikage M, Kamei T, Nakano T, et al. Impact of routine recurrent laryngeal nerve monitoring in prone esophagectomy with mediastinal lymph node dissection. *Surg Endosc*. Jul 2017; 31(7): 2986-2996. PMID 27826777
26. Zhong D, Zhou Y, Li Y, et al. Intraoperative recurrent laryngeal nerve monitoring: a useful method for patients with esophageal cancer. *Dis Esophagus*. Jul 2014; 27(5): 444-51. PMID 23020300
27. Kneist W, Kauff DW, Juhre V, et al. Is intraoperative neuromonitoring associated with better functional outcome in patients undergoing open TME? Results of a case-control study. *Eur J Surg Oncol*. Sep 2013; 39(9): 994-9. PMID 23810330

28. Kneist W, Kauff DW, Rubenwolf P, et al. Intraoperative monitoring of bladder and internal anal sphincter innervation: a predictor of erectile function following low anterior rectal resection for rectal cancer? Results of a prospective clinical study. *Dig Surg.* 2013; 30(4-6): 459-65. PMID 24481247
29. Clarkson JHW, Ozyurekdoglu T, Mujadzic M, et al. An evaluation of the information gained from the use of intraoperative nerve recording in the management of suspected brachial plexus root avulsion. *Plast Reconstr Surg.* Mar 2011; 127(3): 1237-1243. PMID 21364425
30. Zhang W, Chen M, Zhang W, et al. Use of electrophysiological monitoring in selective rhizotomy treating glossopharyngeal neuralgia. *J Craniomaxillofac Surg.* Jul 2014; 42(5): e182-5. PMID 24095216
31. Ochs BC, Herzka A, Y aylali I. Intraoperative neurophysiological monitoring of somatosensory evoked potentials during hip arthroscopy surgery. *Neurodiagn J.* Dec 2012; 52(4): 312-9. PMID 23301281
32. Jahangiri FR. Multimodality neurophysiological monitoring during tibial/fibular osteotomies for preventing peripheral nerve injuries. *Neurodiagn J.* Jun 2013; 53(2): 153-68. PMID 23833842
33. Nagda SH, Rogers KJ, Sestokas AK, et al. Neer Award 2005: Peripheral nerve function during shoulder arthroplasty using intraoperative nerve monitoring. *J Shoulder Elbow Surg.* 2007; 16(3 Suppl): S2-8. PMID 17493556
34. Reddy RP, Chang R, Coutinho DV, et al. Triggered Electromyography is a Useful Intraoperative Adjunct to Predict Postoperative Neurological Deficit Following Lumbar Pedicle Screw Instrumentation. *Global Spine J.* Jun 2022; 12(5): 1003-1011. PMID 34013769
35. Thirumala PD, Crammond DJ, Loke YK, et al. Diagnostic accuracy of motor evoked potentials to detect neurological deficit during idiopathic scoliosis correction: a systematic review. *J Neurosurg Spine.* Mar 2017; 26(3): 374-383. PMID 27935448
36. Alemo S, Sayadipour A. Role of intraoperative neurophysiologic monitoring in lumbosacral spine fusion and instrumentation: a retrospective study. *World Neurosurg.* Jan 2010; 73(1): 72-6; discussion e7. PMID 20452872
37. Bindal RK, Ghosh S. Intraoperative electromyography monitoring in minimally invasive transforaminal lumbar interbody fusion. *J Neurosurg Spine.* Feb 2007; 6(2): 126-32. PMID 17330579
38. Bose B, Wierzbowski LR, Sestokas AK. Neurophysiologic monitoring of spinal nerve root function during instrumented posterior lumbar spine surgery. *Spine (Phila Pa 1976).* Jul 01 2002; 27(13): 1444-50. PMID 12131744
39. Clements DH, Morledge DE, Martin WH, et al. Evoked and spontaneous electromyography to evaluate lumbosacral pedicle screw placement. *Spine (Phila Pa 1976).* Mar 01 1996; 21(5): 600-4. PMID 8852316
40. Darden BV, Wood KE, Hatley MK, et al. Evaluation of pedicle screw insertion monitored by intraoperative evoked electromyography. *J Spinal Disord.* Feb 1996; 9(1): 8-16. PMID 8727451
41. Luo W, Zhang F, Liu T, et al. Minimally invasive transforaminal lumbar interbody fusion aided with computer-assisted spinal navigation system combined with electromyography monitoring. *Chin Med J (Engl).* Nov 2012; 125(22): 3947-51. PMID 23158122
42. Maguire J, Wallace S, Madiga R, et al. Evaluation of intrapedicular screw position using intraoperative evoked electromyography. *Spine (Phila Pa 1976).* May 01 1995; 20(9): 1068-74. PMID 7631237
43. Papadopoulos EC, Girardi FP, Sama A, et al. Accuracy of single-time, multilevel registration in image-guided spinal surgery. *Spine J.* 2005; 5(3): 263-7; discussion 268. PMID 15863081
44. Sutter MA, Eggspuehler A, Grob D, et al. Multimodal intraoperative monitoring (MIOM) during 409 lumbosacral surgical procedures in 409 patients. *Eur Spine J.* Nov 2007; 16 Suppl 2(Suppl 2): S221-8. PMID 17912559
45. Welch WC, Rose RD, Balzer JR, et al. Evaluation with evoked and spontaneous electromyography during lumbar instrumentation: a prospective study. *J Neurosurg.* Sep 1997; 87(3): 397-402. PMID 9285605
46. Wood MJ, Mannion RJ. Improving accuracy and reducing radiation exposure in minimally invasive lumbar interbody fusion. *J Neurosurg Spine.* May 2010; 12(5): 533-9. PMID 20433301
47. Wood MJ, McMillen J. The surgical learning curve and accuracy of minimally invasive lumbar pedicle screw placement using CT based computer-assisted navigation plus continuous electromyography monitoring - a retrospective review of 627 screws in 150 patients. *Int J Spine Surg.* 2014; 8. PMID 25694919
48. Melachuri SR, Melachuri MK, Anetakis K, et al. Diagnostic Accuracy of Thresholds Less Than or Equal to 8mA in Pedicle Screw Testing During Lumbar Spine Procedures to Predict New Postoperative Lower Extremity Neurological Deficits. *Spine (Phila Pa 1976).* Jan 15 2021; 46(2): E139-E145. PMID 33347093
49. Accadbled F, Henry P, de Gauzy JS, et al. Spinal cord monitoring in scoliosis surgery using an epidural electrode. Results of a prospective, consecutive series of 191 cases. *Spine (Phila Pa 1976).* Oct 15 2006; 31(22): 2614-23. PMID 17047554
50. Eggspuehler A, Sutter MA, Grob D, et al. Multimodal intraoperative monitoring during surgery of spinal deformities in 217 patients. *Eur Spine J.* Nov 2007; 16 Suppl 2(Suppl 2): S188-96. PMID 17632737
51. El-Hawary R, Sucato DJ, Sparagana S, et al. Spinal cord monitoring in patients with spinal deformity and neural axis abnormalities: a comparison with adolescent idiopathic scoliosis patients. *Spine (Phila Pa 1976).* Sep 01 2006; 31(19): E698-706. PMID 16946643
52. Feng B, Qiu G, Shen J, et al. Impact of multimodal intraoperative monitoring during surgery for spine deformity and potential risk factors for neurological monitoring changes. *J Spinal Disord Tech.* Jun 2012; 25(4): E108-14. PMID 22367467
53. Kundnani VK, Zhu L, Tak H, et al. Multimodal intraoperative neuromonitoring in corrective surgery for adolescent idiopathic scoliosis: Evaluation of 354 consecutive cases. *Indian J Orthop.* Jan 2010; 44(1): 64-72. PMID 20165679
54. Lo YL, Dan YF, Teo A, et al. The value of bilateral ipsilateral and contralateral motor evoked potential monitoring in scoliosis surgery. *Eur Spine J.* Sep 2008; 17 Suppl 2(Suppl 2): S236-8. PMID 17874145
55. Luk KD, Hu Y, Wong YW, et al. Evaluation of various evoked potential techniques for spinal cord monitoring during scoliosis surgery. *Spine (Phila Pa 1976).* Aug 15 2001; 26(16): 1772-7. PMID 11493849
56. Macdonald DB, Al Zayed Z, Al Saddigi A. Four-limb muscle motor evoked potential and optimized somatosensory evoked potential monitoring with decussation assessment: results in 206 thoracolumbar spine surgeries. *Eur Spine J.* Nov 2007; 16 Suppl 2(Suppl 2): S171-87. PMID 17638028

57. Noonan KJ, Walker T, Feinberg JR, et al. Factors related to false- versus true-positive neuromonitoring changes in adolescent idiopathic scoliosis surgery. *Spine (Phila Pa 1976)*. Apr 15 2002; 27(8): 825-30. PMID 11935104
58. Pastorelli F, Di Silvestre M, Plasmati R, et al. The prevention of neural complications in the surgical treatment of scoliosis: the role of the neurophysiological intraoperative monitoring. *Eur Spine J*. May 2011; 20 Suppl 1(Suppl 1): S105-14. PMID 21416379
59. Pron Y, Bernard JM, Fayet G, et al. Usefulness of neurogenic motor evoked potentials for spinal cord monitoring: findings in 112 consecutive patients undergoing surgery for spinal deformity. *Electroencephalogr Clin Neurophysiol*. Jan 1998; 108(1): 17-23. PMID 9474058
60. Schwartz DM, Auerbach JD, Dormans JP, et al. Neurophysiological detection of impending spinal cord injury during scoliosis surgery. *J Bone Joint Surg Am*. Nov 2007; 89(11): 2440-9. PMID 17974887
61. Murphy B, Murphy E, Irwin J, et al. Review of the Role of Intraoperative Neurophysiological Monitoring in Spinal Surgery With a Focus on the True and False Positives and a Clinical Correlation. *Int J Spine Surg*. Jun 2022; 16(3): 548-553. PMID 35772981
62. Tsirikos AI, Duckworth AD, Henderson LE, et al. Multimodal Intraoperative Spinal Cord Monitoring during Spinal Deformity Surgery: Efficacy, Diagnostic Characteristics, and Algorithm Development. *Med Princ Pract*. 2020; 29(1): 6-17. PMID 31158841
63. American Academy of Neurology. Model Coverage Policy: Principles of Coding for Intraoperative Neurophysiologic Monitoring (IOM) and Testing. 2012; https://www.aan.com/siteassets/home-page/tools-and-resources/practicing-neurologist--administrators/billing-and-coding/model-coverage-policies/16iommodelpolicy_tr.pdf. Accessed March 1, 2023.
64. American Association of Neurological Surgeons (AANS)/Congress of Neurological Surgeons (CNS). Joint Section on Disorders of the Spine and Peripheral Nerves updated position statement: intraoperative electrophysiological monitoring. January 2018. <https://www.spinesection.org/statement-detail/intraoperative-electrophysiological-monitoring>. Accessed March 1, 2023.
65. Sharan A, Groff MW, Dailey AT, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 15: electrophysiological monitoring and lumbar fusion. *J Neurosurg Spine*. Jul 2014; 21(1): 102-5. PMID 24980592
66. American Clinical Neurophysiology Society. Guideline 11A: Recommended Standards for Neurophysiologic Intraoperative Monitoring Principles. 2009; <https://www.acns.org/pdf/guidelines/Guideline-11A.pdf>. Accessed March 3, 2023.
67. Diercks GR, Rastatter JC, Kazahaya K, et al. Pediatric intraoperative nerve monitoring during thyroid surgery: A review from the American Head and Neck Society Endocrine Surgery Section and the International Neural Monitoring Study Group. *Head Neck*. Jun 2022; 44(6): 1468-1480. PMID 35261110
68. Macdonald DB, Skinner S, Shils J, et al. Intraoperative motor evoked potential monitoring - a position statement by the American Society of Neurophysiological Monitoring. *Clin Neurophysiol*. Dec 2013; 124(12): 2291-316. PMID 24055297
69. Halsey MF, Myung KS, Ghag A, et al. Neurophysiological monitoring of spinal cord function during spinal deformity surgery: 2020 SRS neuromonitoring information statement. *Spine Deform*. Aug 2020; 8(4): 591-596. PMID 32451978
70. National Institute for Health and Care Excellence (NICE). Intraoperative nerve monitoring during thyroid surgery [IPG255]. 2008; <https://www.nice.org.uk/guidance/ipg255/chapter/1-guidance>. Accessed March 6, 2023.
71. Centers for Medicare & Medicaid Services. National Coverage Determination (NCD) for Electroencephalographic Monitoring During Surgical Procedures Involving the Cerebral Vasculature (160.8). 2006; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=77&ncdver=2&CoverageSelection=National&Keyword=monitoring&KeywordLookUp=Title&KeywordLookUp=Title&KeywordLookUp=Title&KeywordSearchType=And&KeywordSearchType=And&KeywordSearchType=And&bc=gAAACAAAAA&>. Accessed March 6, 2023.
72. Centers for Medicare & Medicaid Services. Billing Medicare for Remote Intraoperative Neurophysiology Monitoring in CY 2013. Updated September 2020; <https://www.cms.gov/medicare/medicare-fee-for-service-payment/physicianfeesched/downloads/faq-remote-ionm.pdf>. Accessed March 6, 2023.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
December 2011	New policy	
March 2013	Replace policy	Policy updated with literature review; references added and reordered; policy statements unchanged.
September 2014	Replace policy	Policy updated with literature review, references 10-14, 16-18, 22, 24, and 25 added; policy statements unchanged.
September 2015	Replace policy	Policy updated with literature review; references 12, 13, 15, and 22 added; policy statements unchanged.
June 2017	Replace policy	Policy updated with literature review through October 11, 2016; references added and some references removed. Intraoperative monitoring is considered medically necessary for high risk thyroid and anterior cervical spine surgeries. Title changed to "Intraoperative Neurophysiologic Monitoring.,
June 2018	Replace policy	Policy updated with literature review through February 23, 2018; references 8, 10, and 14 added; references 6-7 updated. Policy statements unchanged.
June 2019	Replace policy	Policy updated with literature review through February 18, 2019; references added. Policy statements unchanged.
June 2020	Replace policy	Policy updated with literature review through February 11, 2020; references added. Policy statements unchanged.
June 2021	Replace policy	Policy updated with literature review through March 2, 2021; no references added. Policy statements unchanged.
June 2022	Replace policy	Policy updated with literature review through March 3, 2022; no references added. Policy statement on intraoperative electromyography and nerve conduction velocity monitoring during surgery on the peripheral nerves changed from "not medically necessary" to "investigational"; intent unchanged. Policy statements otherwise unchanged.
June 2023	Replace policy	Policy updated with literature review through March 6, 2023; references added. New indication for spinal instrumentation requiring screws or distraction added. Minor editorial refinements to policy statements; intent unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.